

Access this article online

Quick Response Code:



Website:
https://eurasianj pulmonol.org

DOI:
10.14744/ejp.2024.93364

Sleep-related breathing disorders in idiopathic pulmonary fibrosis: A new hidden perspective

Safy Kaddah¹, Antonio Esquinas²

ORCID:

Safy Kaddah: 0000-0003-3235-4228

Antonio M. Esquinas: 0000-0003-0571-2050

Dear Editor,

We have read the original article entitled "The spectrum of sleep-related breathing disorders (SRBDS) in idiopathic pulmonary fibrosis (IPF)" by Vayisoğlu Şahin et al.^[1] and would like to congratulate the authors for this original analysis. However, we believe that certain aspects require further clarification for their clinical implications.

point could provide more valuable insight for clinical practice.^[2]

Second, the results stated that the mean sleep efficiency percentage of the patients was $66.8 \pm 15.2\%$. A sleep efficiency below 85% is considered poor and may indicate coexisting insomnia. This finding raises the possibility of coexisting insomnia and sleep apnea (COMISA) in these patients, which should be considered.^[3]

First, the authors reported that the mean STOP-Bang scores (Snoring, Tiredness, Observed apnea, high blood Pressure, Body mass index, Age, Neck circumference, and Gender) differed significantly between the two groups: patients with SRBD scored 4.2 ± 1.4 , while those without SRBD scored 3.2 ± 1.1 ($p=0.032$). However, STOP-Bang scores between 3 and 4 indicate an intermediate risk for obstructive sleep apnea (OSA), which places both groups in the same risk category for developing OSA. Clarification of this

Third, the difference in OSA symptoms, specifically apnea and daytime sleepiness, is a key issue. Table 4 presents more than one remarkable finding: patients with SRBD had apnea and daytime sleepiness rates of 4.7% and 23.3%, respectively, compared to 7.7% and 30.8% in patients without SRBD. This suggests that IPF patients without SRBD may experience apnea and daytime sleepiness more frequently than those with SRBD. Additionally, a comparison of the Epworth Sleepiness Scale between the two groups showed similar

¹Department of Pulmonology, Faculty of Medicine, Kasr Alainy Hospital, Cairo University, Cairo, Egypt,
²Intensive Care Unit, Hospital Meseguer, NIV-ICM, Biomedical Research Institute of Murcia Pascual Parrilla-IMIB, Murcia, Spain

Address for correspondence:

Dr. Safy Kaddah,
Department of Pulmonology, Faculty of Medicine, Kasr Alainy Hospital, Cairo University, Cairo, Egypt.
E-mail: safykaddah@kasralainy.edu.eg

Received: 09-09-2024
Accepted: 17-09-2024
Published: 27-06-2025

How to cite this article: Kaddah S, Esquinas A. Sleep-related breathing disorders in idiopathic pulmonary fibrosis: A new hidden perspective. Eurasian J Pulmonol 0000;00:1-2.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: kare@karepb.com



results: 5.8 ± 4.6 for patients with SRBD versus 5.6 ± 5 for those without SRBD. Furthermore, the Apnea-Hypopnea Index (AHI) was found to be 19.2 ± 19.3 in IPF patients with SRBD and 6.7 ± 10.5 in those without SRBD. It was previously mentioned in the results section on page 104 that “The mean AHI was 15.1 ± 17.3 .^[4] An AHI value above 5 was observed in 43 patients.” Based on these combined findings, we considered the possibility that a significant proportion of the 55 IPF patients enrolled in the study could be diagnosed with SRBD.^[5]

Fourth, it is important to consider the overlapping nature of symptoms such as dyspnea and cough symptoms in the categorical analysis of IPF symptoms. Although statistical significance was observed regarding IPF symptoms between the two groups, we noticed that cough—either alone or in combination with other symptoms—appeared to be a major factor interfering with the sleep quality of IPF patients. This effect was especially pronounced in the group with SRBDs, intensifying their complaints.^[6]

Finally, it is important to consider the evaluation of oxygenation parameters. It would provide valuable information if the authors could provide results for minimum O_2 saturation (%), average O_2 saturation (%), O_2 saturation over 90% (%), oxygen desaturation index (ODI), and average O_2 saturation during wakefulness (%) between IPF patients receiving oxygen therapy and those who are not. This data could offer further observational insight into whether the need for oxygen therapy influences OSA symptoms and sleep quality.^[7]

Further clinical studies are needed to clarify the spectrum of SRBDs in IPF and their implications for early diagnosis.

Conflicts of Interest Statement

The authors have no conflicts of interest to declare.

Funding

The authors declared that this study received no financial support.

Use of AI for Writing Assistance

No AI technologies utilized.

Authorship Contributions

Concept – S.K.; Design – S.K.; Supervision – S.K.; Funding – S.K.; Analysis and/or interpretation – S.K., A.E.; Literature search – A.E.; Writing – S.K.; Critical review – S.K., A.E.

Peer-review

Externally peer-reviewed.

References

1. Vayisoğlu Şahin G, Yalnız Ö, Polat G, Yalnız E, Demirci Üçsular F, Uçar ZZ. The spectrum of sleep-related breathing disorders in idiopathic pulmonary fibrosis. *Eurasian J Pulmonol* 2024;26(2):102–11. [\[CrossRef\]](#)
2. Chung F, Yegneswaran B, Liao P, Chung SA, Vairavanathan S, Islam S, et al. STOP questionnaire: a tool to screen patients for obstructive sleep apnea. *Anesthesiology* 2008;108(5):812–21. [\[CrossRef\]](#)
3. Wulterkens BM, Hermans LWA, Fonseca P, Asin J, Duis N, Janssen HCJP, et al. Sleep structure in patients with COMISA compared to OSA and insomnia. *J Clin Sleep Med* 2023;19(6):1051–9. [\[CrossRef\]](#)
4. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991;14(6):540–5. [\[CrossRef\]](#)
5. Malhotra A, Ayappa I, Ayas N, Collop N, Kirsch D, Mcardle N, et al. Metrics of sleep apnea severity: beyond the apnea-hypopnea index. *Sleep* 2021;44(7):zsab030. [\[CrossRef\]](#)
6. Pfeiffer M. Restriktive Lungenerkrankungen. In: *Enzyklopädie der Schlafmedizin*. Peter H, Penzel T, Peter JH, Peter JG, editors. Heidelberg: Springer; 2007. p.1026–31.
7. Kolilekas L, Manali E, Vlami KA, Lyberopoulos P, Triantafyllidou C, Kagouridis K, et al. Sleep oxygen desaturation predicts survival in idiopathic pulmonary fibrosis. *J Clin Sleep Med* 2013;9(6):593–601. [\[CrossRef\]](#)